arXiv:0704.3523v1 [math.AG] 26 Apr 2007

Immersions of spheres and algebraically constructible functions

Iwona Karolkiewicz Aleksandra Nowel
Zbigniew Szafraniec

April 2007

Abstract

Let Λ be an algebraic set and let $g : \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R}^{2n}$ (n is even) be a polynomial mapping such that for each $\lambda \in \Lambda$ there is $r(\lambda) > 0$ such that the mapping $g_\lambda = g(\cdot, \lambda)$ restricted to the sphere $S^n(r)$ is an immersion for every $0 < r < r(\lambda)$, so that the intersection number $\mathrm{I}(g_\lambda | S^n(r))$ is defined. Then $\Lambda \ni \lambda \mapsto \mathrm{I}(g_\lambda | S^n(r)) \in \mathbb{Z}$ is an algebraically constructible function.

Keywords: immersions of spheres, algebraically constructible functions, real algebraic sets.
Mathematics Subject Classification (2000): 14P25 57N35 32S50

1 Introduction

McCrory and Parusiński [6] have introduced algebraically constructible functions in order to study the topology of real algebraic sets.

Let Λ be a real algebraic set. An integer valued function $\phi : \Lambda \longrightarrow \mathbb{Z}$ is *algebraically constructible* if there exist an algebraic set W and a proper regular map $p : W \longrightarrow \Lambda$ such that $\phi(\lambda)$ equals the Euler characteristic $\chi(p^{-1}(\lambda))$.

If that is the case then ϕ is *semialgebraically constructible*, that is there exists a semialgebraic stratification $\mathcal{S}$ of Λ such that ϕ is constant on strata of

---

Iwona Karolkiewicz, Aleksandra Nowel and Zbigniew Szafraniec
University of Gdańsk, Institute of Mathematics
80-952 Gdańsk, Wita Stwosza 57
Poland
Tel.: +48-58-5232059
Fax: +48-58-3414914
Email: ikarolki@manta.univ.gda.pl
Email: Aleksandra.Nowel@math.univ.gda.pl
Email: Zbigniew.Szafraniec@math.univ.gda.pl

First and second named authors supported by the grant BW /5100-5-0286-7

$\mathcal{S}$. If Λ is irreducible then ϕ has to be generically constant modulo 2, see for instance [1, Proposition 2.3.2], and there exist a real polynomial $g : \Lambda \longrightarrow \mathbb{R}$ and a constant μ such that generically on Λ: $\phi \equiv \mu + \operatorname{sgn} g \mod 4$ [3].

In fact, algebraically constructible functions are precisely those constructible functions which are sums of signs of polynomials [9], [10].

Let $f : \mathbb{R}^n \times \Lambda \longrightarrow \mathbb{R}^n$ be a polynomial mapping such that $0 \in \mathbb{R}^n$ is isolated in $f(\cdot, \lambda)^{-1}(0)$ for all $\lambda \in \Lambda$, so that the local topological degree $\deg_0 f(\cdot, \lambda)$ of $f(\cdot, \lambda)$ at the origin is well defined. Then $\Lambda \ni \lambda \mapsto \deg_0 f(\cdot, \lambda) \in \mathbb{Z}$ is algebraically constructible [9, Theorem 3.4].

For a recent account of the theory we refer the reader to [2], [7], [8].

Whitney [13] has introduced an intersection number $\mathrm{I}(g)$ for an immersion $g : M^n \longrightarrow \mathbb{R}^{2n}$. If n is even then $\mathrm{I}(g) \in \mathbb{Z}$, if n is odd then $\mathrm{I}(g) \in \mathbb{Z}_2$. Smale [11] proved, that two immersions $f, g : S^n \longrightarrow \mathbb{R}^{2n}$ are regularly homotopic if and only if $\mathrm{I}(f) = \mathrm{I}(g)$.

In this paper we investigate how does the "local" intersection number change when there is an algebraic family of immersions. Let $g = (g_1, \ldots, g_{2n}) : \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R}^{2n}$ (n is even) be a polynomial mapping. Assume that for each $\lambda \in \Lambda$ there exists $r(\lambda) > 0$ such that the mapping $g_\lambda = g(\cdot, \lambda)$ restricted to the sphere $S^n(r)$ is an immersion for $0 < r < r(\lambda)$. If that is the case then the intersection number $\mathrm{I}(g_\lambda | S^n(r))$ is the same for all $0 < r < r(\lambda)$. We shall prove (Theorem 6.1) that the function

$$\Lambda \ni \lambda \mapsto \mathrm{I}(g_\lambda | S^n(r)) \in \mathbb{Z}$$

is algebraically constructible.

2 Preliminaries

Suppose that L is a p–dimensional oriented manifold, $H : L \longrightarrow \mathbb{R}^p$ is a smooth mapping, and U is an open subset of L such that $H^{-1}(0) \cap U$ is compact.

There exists $(N, \partial N)$ — a compact p–dimensional oriented manifold with boundary such that $N \subset U$ and $H^{-1}(0) \cap U \subset N \setminus \partial N$.

By *the topological degree* of the mapping

$$(U, U \setminus H^{-1}(0)) \ni x \mapsto H(x) \in (\mathbb{R}^p, \mathbb{R}^p \setminus \{0\})$$

we mean the topological degree of

$$(N, \partial N) \ni x \mapsto H(x) \in (\mathbb{R}^p, \mathbb{R}^p \setminus \{0\})$$

which equals the degree of the mapping

$$\partial N \ni x \mapsto \frac{H(x)}{\|H(x)\|} \in S^{p-1}.$$

Of course, the degree does not depend on the choice of N.

Let $F_1, \ldots, F_k; G_1, \ldots, G_{n-k} : \mathbb{R}^n, 0 \to \mathbb{R}, 0$ be analytic functions defined in a neighbourhood of the origin.

Denote

$$\begin{aligned} F &= (F_1, \ldots, F_k) : \mathbb{R}^n, 0 \to \mathbb{R}^k, 0 \\ G &= (G_1, \ldots, G_{n-k}) : \mathbb{R}^n, 0 \to \mathbb{R}^{n-k}, 0 \\ S^{n-1}(r) &= \{x \in \mathbb{R}^n \mid \|x\| = r\} \\ S^{n-1} &= \{x \in \mathbb{R}^n \mid \|x\| = 1\} \\ B^n(r) &= \{x \in \mathbb{R}^n \mid \|x\| \leq r\} \end{aligned}$$

Suppose that $F^{-1}(0)$ has an isolated singularity at the origin, i.e. for $\|x\|$ small enough, if $F(x) = 0$ and $\operatorname{rank}[DF(x)] < k$ then $x = 0$. If $r > 0$ is small enough then $S^{n-1}(r)$ cuts $F^{-1}(0)$ transversally, so $M(r) = S^{n-1}(r) \cap F^{-1}(0)$ is either void or a compact $(n-k-1)$–dimensional manifold.

We shall say that vectors $v_1, \ldots, v_{n-k-1}$ in the tangent space $T_xM(r)$ are *well oriented* if $\nabla F_1(x), \ldots, \nabla F_k(x), x, v_1, \ldots, v_{n-k-1}$ are well oriented in $\mathbb{R}^n$. This way $M(r)$ is oriented.

Let $y \in \mathbb{R}^k$ be a regular value of F. Then $F^{-1}(y)$ is either void or an $(n-k)$–dimensional manifold. We shall say that $w_1, \ldots, w_{n-k} \in T_xF^{-1}(y)$ are *well oriented* if $\nabla F_1(x), \ldots, \nabla F_k(x), w_1, \ldots, w_{n-k}$ are well oriented in $\mathbb{R}^n$. This way $F^{-1}(y)$ is also oriented.

Fix small $r > 0$. If y lies sufficiently close to the origin then $S^{n-1}(r)$ cuts $F^{-1}(y)$ transversally, so $\overline{M}(r) = S^{n-1}(r) \cap F^{-1}(y)$ is either void or a compact $(n-k-1)$–dimensional manifold. Moreover, $B^n(r) \cap F^{-1}(y)$ is a compact oriented manifold with boundary $\partial\left(B^n(r) \cap F^{-1}(y)\right) = \overline{M}(r)$. The orientation of the boundary induced from $B^n(r) \cap F^{-1}(y)$ may be described the same way as the orientation of $M(r)$. Manifolds $M(r)$ and $\overline{M}(r)$ are isotopic.

Suppose that $F^{-1}(0) \cap G^{-1}(0) = \{0\}$. Then $M(r) \cap G^{-1}(0) = \emptyset$ and $\overline{M}(r) \cap G^{-1}(0) = \emptyset$.

If $\|y\|$ is small enough then mappings

$$M(x) \ni x \mapsto \frac{G(x)}{\|G(x)\|} \in S^{n-k-1},$$

$$\overline{M}(x) \ni x \mapsto \frac{G(x)}{\|G(x)\|} \in S^{n-k-1}$$

have the same topological degree. Denote it by $\deg(M(r))$. It equals the degree ρ of $\overline{G}$, where $\overline{G}$ is the restricted mapping

$$(F^{-1}(y) \cap B^n(r), \overline{M}(r)) \ni x \mapsto G(x) \in (\mathbb{R}^{n-k}, \mathbb{R}^{n-k} \setminus \{0\}).$$

Choose a regular value $z \in \mathbb{R}^{n-k}$ near the origin. Then

$$\rho = \sum \operatorname{sgn} \det[D\,\overline{G}(x)] \qquad (\text{where } x \in \overline{G}^{-1}(z)).$$

Using the concept of the Gram determinant the reader may check that

$$\rho = \sum \operatorname{sgn} \det[D(F,G)(x)],$$

where $x \in F^{-1}(y) \cap B^n(r) \cap G^{-1}(z) = (F,G)^{-1}(y,z) \cap B^n(r)$. If $0 < \|z\| \ll \|y\| \ll r \ll 1$ then ρ equals the local topological degree $\deg_0(F,G)$ at the origin, i.e. the topological degree of the mapping

$$S^{n-1}(r) \ni x \mapsto \frac{(F(x),G(x))}{\|(F(x),G(x))\|} \in S^{n-1}.$$

Set

$$N(r) = \{x \in S^{n-1}(r) \mid F_1(x) = \ldots = F_{k-1}(x) = 0, F_k(x) \geq 0\}.$$

Suppose that $(F_1,\ldots,F_{k-1})^{-1}(0)$ has an isolated singularity at the origin. Since $F^{-1}(0)$ has an isolated singularity, $N(r)$ is either void or an $(n-k)$–dimensional compact oriented manifold with boundary, and $\partial N(r) = S^{n-1}(r) \cap F^{-1}(0) = M(r)$. There is the restricted function

$$(N(r), \partial N(r)) \ni x \mapsto G(x) \in (\mathbb{R}^{n-k}, \mathbb{R}^{n-k} \setminus \{0\}),$$

which will be denoted by $\overline{G}_r$.

The topological degree of $\overline{G}_r$ equals the topological degree of

$$\partial N(r) \ni x \mapsto \frac{G(x)}{\|G(x)\|} \in S^{n-k-1},$$

which is the same as $\deg(M(r)) = \rho = \deg_0(F,G)$. We have proved:

Proposition 2.1 *Let $F_1,\ldots,F_k; G_1,\ldots,G_{n-k} : \mathbb{R}^n, 0 \to \mathbb{R}, 0$ be analytic functions defined in a neighbourhood of the origin. Suppose that $(F_1,\ldots,F_{k-1})^{-1}(0)$, as well as $(F_1,\ldots,F_k)^{-1}(0)$, has an isolated singularity at the origin, and $\{0\}$ is isolated in $(F_1,\ldots,F_k; G_1,\ldots,G_{n-k})^{-1}(0)$.*

If $r > 0$ is small enough then the topological degree of the mapping $\overline{G}_r$, i.e. of

$$(N(r), \partial N(r)) \ni x \mapsto G(x) \in (\mathbb{R}^{n-k}, \mathbb{R}^{n-k} \setminus \{0\}),$$

does not depend on r and is equal to the local topological degree $\deg_0(F,G)$.

□

Let $\delta : \mathbb{R}^n, 0 \to \mathbb{R}, 0$ be an analytic function. In [12] there is proven that if α is a sufficiently large positive even integer and $t \neq 0$ then $\delta - t\|x\|^\alpha$ has an isolated critical point at the origin. Using the same arguments one may prove:

Proposition 2.2 *Suppose that $(F_1,\ldots,F_{k-1})^{-1}(0)$ has an isolated singularity at the origin.*

Then there exists $\alpha_0 > 0$ such that for any even integer $\alpha > \alpha_0$ and $t \neq 0$

$$(F_1,\ldots,F_{k-1}, \delta - t\|x\|^\alpha)^{-1}(0)$$

has an isolated singularity at the origin.

□

Suppose that $\delta \geq 0$ and

$$X := \{0\} \cup ((F_1, \dots, F_{k-1}, G_1, \dots, G_{n-k})^{-1}(0) \setminus \delta^{-1}(0))$$

is closed. Then $\delta^{-1}(0) \cap X = \{0\}$. Since X is closed semianalytic, there exists a Łojasiewicz exponent α_1, such that $\delta(x) > t\|x\|^{\alpha_1}$ for $t > 0$ and $x \in X \setminus \{0\}$ sufficiently close to the origin.

Set

$$L(r) = \{x \in S^{n-1}(r) \mid F_1(x) = \ldots = F_{k-1}(x) = 0\}.$$

Then $U(r) = L(r) \setminus \delta^{-1}(0)$ is an open subset of $L(r)$, and $U(r) \cap G^{-1}(0) = S^{n-1}(r) \cap X$ is a compact subset of $U(r)$.

Let $\alpha > \max(\alpha_0, \alpha_1)$ be an even positive integer. Set $F_k = \delta - t\|x\|^{\alpha}$ for arbitrary $t > 0$. Then $F_k(x) > 0$ for $x \in L(r) \cap G^{-1}(0) \setminus \delta^{-1}(0)$, and $F_k(x) < 0$ for $x \in L(r) \cap G^{-1}(0) \cap \delta^{-1}(0)$.

Proposition 2.3 *Suppose that*

(a) $(F_1, \dots, F_{k-1})^{-1}(0)$ *has an isolated singularity at the origin, so that* $L(r)$ *is either void or a compact oriented* $(n-k)$*–dimensional manifold for small* $r > 0$,

(b) $\delta \geq 0$ *and* $U(r) \cap G^{-1}(0) = L(r) \cap G^{-1}(0) \setminus \delta^{-1}(0)$ *is a compact subset of* $U(r)$.

Then there is a mapping G_r *given by*

$$(U(r), U(r) \setminus G^{-1}(0)) \ni x \mapsto G(x) \in (\mathbb{R}^{n-k}, \mathbb{R}^{n-k} \setminus \{0\})$$

such that if $\alpha > \max(\alpha_0, \alpha_1)$ *is an even integer,* $t > 0$, *and* $F_k = \delta - t\|x\|^{\alpha}$, *then*

$$\mathbb{R}^n \ni x \mapsto (F(x), G(x)) \in \mathbb{R}^n$$

has an isolated zero at the origin, and for each $r > 0$ *small enough the topological degree of* G_r *equals the local topological degree* $\deg_0(F, G)$.

Proof. Since $F_k(x) > 0$ for $x \in U(r) \cap G^{-1}(0) = L(r) \cap G^{-1}(0) \setminus \delta^{-1}(0)$, then $U(r) \cap G^{-1}(0) \subset N(r) \setminus \partial N(r) \subset U(r)$. So the topological degree of G_r equals the topological degree of $\overline{G}_r$, and by Proposition 2.1, equals $\deg_0(F, G)$. □

3 The intersection number of an immersion

Let M be an n–dimensional manifold. A C^1 map $g : M \longrightarrow \mathbb{R}^m$ is called *an immersion* if for each $p \in M$ the rank of $Dg(p)$ equals n.

A homotopy $h_t : M \longrightarrow \mathbb{R}^m$ is called *a regular homotopy,* if at each stage it is an immersion and the induced homotopy of the tangent bundle is continuous.

Theorem 3.1 [11, Theorem B]) *Two C^∞ immersions from an n–dimensional sphere S^n to $\mathbb{R}^m$ are regularly homotopic when $m \geqslant 2n+1$.*

As in [13] we say that an immersion $g : M \longrightarrow \mathbb{R}^{2n}$ has a *regular self–intersection* at the point $g(p) = g(q)$ if

$$Dg(p)T_pM + Dg(q)T_qM = \mathbb{R}^{2n}.$$

An immersion $g : M \longrightarrow \mathbb{R}^{2n}$ is called *completely regular* if it has only regular self–intersections and no triple points.

Assume that n is **even** and M is compact and oriented.

Let $g : M \longrightarrow \mathbb{R}^{2n}$ be a completely regular immersion and have a regular self–intersection at the point $g(p) = g(q)$.

Let $u_1, \ldots, u_n \in T_pM$, $v_1, \ldots, v_n \in T_qM$ be sets of well–oriented, independent vectors in respective tangent spaces of M. Then the vectors $Dg(p)u_1, \ldots, Dg(p)u_n$, $Dg(q)v_1, \ldots, Dg(q)v_n$ form a basis in $\mathbb{R}^{2n}$. As in [13] we will say that the self–intersection at the point $g(p) = g(q)$ is *positive* or *negative* according to whether this basis determines the positive or negative orientation of $\mathbb{R}^{2n}$.

The intersection number of a completely regular immersion g is the algebraic number of its self–intersections.

Let for a given (not necessarily completely regular) immersion $g : M \longrightarrow \mathbb{R}^{2n}$ define $G : M \times M \longrightarrow \mathbb{R}^{2n}$ as $G(x,y) = g(x) - g(y)$. Set

$$\Delta = \{(p,p) \mid p \in M\} \subset M \times M.$$

Since g is an immersion, Δ is isolated in $G^{-1}(0)$, and so $G^{-1}(0) \setminus \Delta$ is a compact subset of $M \times M \setminus \Delta$. Of course, $M \times M \setminus \Delta$ is an open subset of $M \times M$, and

$$(M \times M \setminus \Delta) \setminus G^{-1}(0) = (M \times M \setminus (G^{-1}(0) \cup \Delta)).$$

The topological degree $d(g)$ of the mapping

$$(M \times M \setminus \Delta, M \times M \setminus (G^{-1}(0) \cup \Delta)) \ni (x,y) \mapsto G(x,y) \in (\mathbb{R}^{2n}, \mathbb{R}^{2n} \setminus \{0\})$$

is always an even integer. Let us denote

$$\mathrm{I}(g) = \frac{1}{2} d(g).$$

By [5, Theorem 3.1] we have

Theorem 3.2 *If g is a completely regular immersion then its intersection number is equal to $\mathrm{I}(g)$.*

Assume that either n is **odd** or M is non–orientable.

In this case one can also define *the intersection number* of a completely regular immersion $g : M \longrightarrow \mathbb{R}^{2n}$ as the number of its self–intersections modulo 2.

By [13, Theorem 2] if M is closed then the intersection number is invariant under regular homotopies. As in [14], if M is closed, any immersion $g : M \longrightarrow \mathbb{R}^{2n}$ can be made completely regular by a regular homotopy. So in this case we can define the intersection number also for g that is not completely regular.

We have a characterisation of regularly homotopic immersions due to Smale [11]:

Theorem 3.3 [11, Theorem C] *Two C^∞ immersions f, g from an n–dimensional sphere S^n to $\mathbb{R}^{2n}$ are regularly homotopic if and only if* $\mathrm{I}(f) = \mathrm{I}(g)$.

4 Immersions on small spheres

Let

$$h = (h_1, \ldots, h_l) : \mathbb{R}^n \longrightarrow \mathbb{R}^l$$

$$g = (g_1, \ldots, g_k) : \mathbb{R}^n \longrightarrow \mathbb{R}^k$$

be C^1 mappings. Put $M := h^{-1}(0)$. Suppose that each point $p \in M$ is a regular point of h, i.e. the rank of the derivative matrix $Dh(p)$:

$$\begin{bmatrix} \frac{\partial h_1}{\partial x_1}(p) & \frac{\partial h_1}{\partial x_2}(p) & \cdots & \frac{\partial h_1}{\partial x_n}(p) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial h_l}{\partial x_1}(p) & \frac{\partial h_l}{\partial x_2}(p) & \cdots & \frac{\partial h_l}{\partial x_n}(p) \end{bmatrix}$$

equals l at each $p \in M$. If that is the case then M is a C^1 $(n - l)$–manifold, and there is the restricted mapping $g|M : M \longrightarrow \mathbb{R}^k$.

Proposition 4.1

$$\mathrm{rank}\left[\ D(g|M)(p)\ \right] = \mathrm{rank} \begin{bmatrix} \frac{\partial g_1}{\partial x_1}(p) & \cdots & \frac{\partial g_1}{\partial x_n}(p) \\ \vdots & \ddots & \vdots \\ \frac{\partial g_k}{\partial x_1}(p) & \cdots & \frac{\partial g_k}{\partial x_n}(p) \\ \frac{\partial h_1}{\partial x_1}(p) & \cdots & \frac{\partial h_1}{\partial x_n}(p) \\ \vdots & \ddots & \vdots \\ \frac{\partial h_l}{\partial x_1}(p) & \cdots & \frac{\partial h_l}{\partial x_n}(p) \end{bmatrix} - l$$

at each point $p \in M$.

Corollary 4.2 *The mapping* $g|M : M \longrightarrow \mathbb{R}^k$ *is an immersion (i.e.* $\operatorname{rank} Dg|M \equiv n - l$*) if and only if at each* $p \in M$

$$\operatorname{rank} \begin{bmatrix} \frac{\partial g_1}{\partial x_1}(p) & \cdots & \frac{\partial g_1}{\partial x_n}(p) \\ \vdots & \ddots & \vdots \\ \frac{\partial g_k}{\partial x_1}(p) & \cdots & \frac{\partial g_k}{\partial x_n}(p) \\ \frac{\partial h_1}{\partial x_1}(p) & \cdots & \frac{\partial h_1}{\partial x_n}(p) \\ \vdots & \ddots & \vdots \\ \frac{\partial h_l}{\partial x_1}(p) & \cdots & \frac{\partial h_l}{\partial x_n}(p) \end{bmatrix} = n,$$

i.e. this matrix has a non-zero $(n \times n)$*–minor.*

Corollary 4.3 *Let* $g = (g_1, g_2, \ldots, g_{2n}) : \mathbb{R}^{n+1}, 0 \longrightarrow \mathbb{R}^{2n}, 0$ *be a* C^1 *function. Let us denote* $\omega(x) := x_1^2 + x_2^2 + \ldots + x_{n+1}^2$ *for* $x \in \mathbb{R}^{n+1}$*. Of course,* $S^n(r) = \{x \mid \omega(x) - r^2 = 0\}$*.*

Then the following conditions are equivalent:

- *(a) there exists* $r_0 > 0$ *such that* g *restricted to each sphere of a radius* $0 < r \leqslant r_0$ *is an immersion;*
- *(b) if* $M_1(x), \ldots, M_N(x)$ *are all the* $(n+1) \times (n+1)$*–minors of the matrix*

$$\begin{bmatrix} \frac{\partial g_1}{\partial x_1} & \frac{\partial g_1}{\partial x_2} & \cdots & \frac{\partial g_1}{\partial x_{n+1}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g_{2n}}{\partial x_1} & \frac{\partial g_{2n}}{\partial x_2} & \cdots & \frac{\partial g_{2n}}{\partial x_{n+1}} \\ \frac{\partial \omega}{\partial x_1} & \frac{\partial \omega}{\partial x_2} & \cdots & \frac{\partial \omega}{\partial x_{n+1}} \end{bmatrix},$$

  then there exists $r_0 > 0$ *such that for each* $x \in \mathbb{R}^{n+1}$ *with* $0 < \|x\| < r_0$ *there exists* $i \in \{1, \ldots, N\}$ *such that* $M_i(x) \neq 0$*.*

Let $g : \mathbb{R}^{n+1} \longrightarrow \mathbb{R}^{2n}$ be a C^2 mapping. Assume that for each $r \in [r_1, r_2]$, where $0 < r_1 < r_2$, $g|S^n(r)$ is an immersion. We can define a regular homotopy

$$H : [0;1] \times S^n(r_1) \longrightarrow \mathbb{R}^{2n},$$

$$(t, x) \mapsto g\left(\left(1 - t + \frac{tr_2}{r_1}\right) x\right)$$

between $H(0, x) = g|S^n(r_1)(x)$ and $H(1, x) = g\left(\frac{r_2}{r_1} x\right)$. By Theorem 3.3, $\mathrm{I}(g|S^n(r_1)) = \mathrm{I}(g(\frac{r_2}{r_1}x)|S^n(r_1)) = \mathrm{I}(g|S^n(r_2))$.

If there exists $r_0 > 0$ such that g restricted to a sphere of a radius $0 < r \leqslant r_0$ is an immersion then the intersection number of each $g|S^n(r)$ is defined and does not depend on r.

Let n be an even positive integer and let

$$g = (g_1, \ldots, g_{2n}) : \mathbb{R}^{n+1} \longrightarrow \mathbb{R}^{2n}$$

be a C^2 mapping. Suppose that g restricted to each sphere of a radius small enough is a completely regular immersion.

Let $s \in \mathbb{R}$, $s \neq 0$, and consider a new mapping $g_s = (sg_1, g_2, \ldots, g_{2n})$. Then g_s restricted to a sphere of a radius small enough is an immersion.

Indeed, let $\tilde{M}_1, \ldots, \tilde{M}_N$ be all the $(n+1) \times (n+1)$–minors of the matrix:

$$\begin{bmatrix} s\frac{\partial g_1}{\partial x_1} & s\frac{\partial g_1}{\partial x_2} & \cdots & s\frac{\partial g_1}{\partial x_{n+1}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g_{2n}}{\partial x_1} & \frac{\partial g_{2n}}{\partial x_2} & \cdots & \frac{\partial g_{2n}}{\partial x_{n+1}} \\ \frac{\partial \omega}{\partial x_1} & \frac{\partial \omega}{\partial x_2} & \cdots & \frac{\partial \omega}{\partial x_{n+1}} \end{bmatrix}.$$

Then either $\tilde{M}_j = sM_j$ or $\tilde{M}_j = M_j$, where M_j is the appropriate minor corresponding to the function g. By Corollary 4.3, there exists $r_0 > 0$ such that for each $0 < \|x\| < r_0$ there exists such i that $M_i(x) \neq 0$, and so $\tilde{M}_i(x) \neq 0$. It means that for each $s \in \mathbb{R} \setminus \{0\}$ the mapping g_s restricted to a sphere of a radius small enough is an immersion.

Let $r > 0$ be small and let $p, q \in S^n(r)$. Then $g(p) = g(q)$ if and only if $g_s(p) = g_s(q)$. Since g restricted to each sphere of a small radius is completely regular, so is g_s.

Corollary 4.4 $\mathrm{I}(g_s|S^n(r)) = \mathrm{sgn}(s)\,\mathrm{I}(g|S^n(r))$.

Proof. Take $p, q \in S^n(r)$ such that $p \neq q$ and $g(p) = g(q)$. Suppose that $v_1, \ldots, v_n$ form a well–oriented basis in $T_pS^n(r)$, and $w_1, \ldots, w_n$ form a well–oriented basis in $T_qS^n(r)$.

For any vector w, the first coordinate of $Dg_s(p)w$ is equal to the first coordinate of $Dg(p)w$ multiplied by s, the other coordinates of $Dg_s(p)w$ are the same as the appropriate coordinates of $Dg(p)w$. Consider two matrices:

$$[Dg_s(p)v_1 \ldots, Dg_s(p)v_n, Dg_s(q)w_1, \ldots, Dg_s(q)w_n]$$

and

$$[Dg(p)v_1 \ldots, Dg(p)v_n, Dg(q)w_1, \ldots, Dg(q)w_n].$$

The first row in the first matrix is equal to the first row in the second matrix muptiplied by s, other respective rows are identical. Then

$$\begin{aligned} &\det[Dg_s(p)v_1 \ldots, Dg_s(p)v_n, Dg_s(q)w_1, \ldots, Dg_s(q)w_n] \\ &= s \cdot \det[Dg(p)v_1 \ldots, Dg(p)v_n, Dg(q)w_1, \ldots, Dg(q)w_n]. \end{aligned}$$

Hence

$$\begin{aligned} &\mathrm{I}(g_s|S^n(r)) \\ &= \frac{1}{2}\sum \mathrm{sgn}(\det[Dg_s(p)v_1 \ldots, Dg_s(p)v_n, Dg_s(q)w_1, \ldots, Dg_s(q)w_n]) \\ &= \mathrm{sgn}(s)\frac{1}{2}\sum \mathrm{sgn}(\det[Dg(p)v_1 \ldots, Dg(p)v_n, Dg(q)w_1, \ldots, Dg(q)w_n]) \\ &= \mathrm{sgn}(s)I(g|S^n(r)), \end{aligned}$$

where $p, q \in S^n(r)$, such that $p \neq q$ and $g_s(p) = g_s(q)$.

$\square$

Hence if g restricted to a sphere of a radius small enough is a completely regular immersion, then we have a family $\{g_s\}$ of C^2 mappings and the intersection number of their restrictions to a sphere of a small radius satisfies:

$$\mathbb{R} \setminus \{0\} \ni s \mapsto \mathrm{I}(g_s|S^n(r)) = \mathrm{sgn}(s)\, \mathrm{I}(g|S^n(r)) \in \mathbb{Z},$$

which is of course determined by a sign of a polynomial. In particular the function $s \mapsto \mathrm{I}(g_s|S^n(r))$ is algebraically constructible.

Now let us consider the mapping

$$g = (g_1, g_2, g_3, g_4) : \mathbb{R}^3 \longrightarrow \mathbb{R}^4$$

$$g(x, y, z) = (x, y, xz, yz).$$

Then for each $r > 0$, $g|S^2(r)$ is an immersion because the matrix:

$$\begin{bmatrix} \frac{\partial g_1}{\partial x_1} & \frac{\partial g_1}{\partial x_2} & \frac{\partial g_1}{\partial x_3} \\ \dots & \dots & \dots \\ \frac{\partial g_4}{\partial x_1} & \frac{\partial g_4}{\partial x_2} & \frac{\partial g_4}{\partial x_3} \\ \frac{\partial \omega}{\partial x_1} & \frac{\partial \omega}{\partial x_2} & \frac{\partial \omega}{\partial x_3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ z & 0 & x \\ 0 & z & y \\ 2x & 2y & 2z \end{bmatrix}$$

has a non-zero (3×3)–minor at each point $p \in \mathbb{R}^3 \setminus \{0\}$. It is easy to verify that $g|S^2(r)(p) = g|S^2(r)(q)$ if and only if $p = (0, 0, r)$ and $q = (0, 0, -r)$.

At $p = (0, 0, r)$

$$Dg(p) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ r & 0 & 0 \\ 0 & r & 0 \end{bmatrix}$$

and vectors $v_1 = (1, 0, 0)$ and $v_2 = (0, 1, 0)$ form a well-oriented basis in $T_pS^2(r)$.

At $q = (0, 0, -r)$

$$Dg(q) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -r & 0 & 0 \\ 0 & -r & 0 \end{bmatrix}$$

and vectors $w_1 = (1, 0, 0)$ and $w_2 = (0, -1, 0)$ form a well-oriented basis in $T_qS^2(r)$.

Since

$$\det[Dg(p)v_1, Dg(p)v_2, Dg(q)w_1, Dg(q)w_2] = \det \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 \\ r & 0 & -r & 0 \\ 0 & r & 0 & r \end{bmatrix} = -4r^2,$$

$g|S^2(r)$ is a completely regular immersion and

$$\mathrm{I}(g|S^2(r)) = -1.$$

Then for each $s \in \mathbb{R}\setminus\{0\}$, the mapping $g_s = (sx, y, xz, yz)$ restricted to $S^2(r)$ is a completely regular immersion, and

$$\mathrm{I}(g_s|S^2(r)) = -\operatorname{sgn}(s).$$

So $\mathbb{R}\setminus\{0\} \ni s \mapsto \mathrm{I}(g_s|S^2(r)) = -\operatorname{sgn} s \in \{-1, 1\}$ is an algebraically constuctible function, which is of course nontrivial and constant modulo 2, but not constant modulo 4.

5 Families of analytic mappings

Let $\Lambda \subset \mathbb{R}^p$ be an analytic set. Let

$$g = (g_1, \ldots, g_{2n}) : \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R}^{2n},$$

where $g_1, \ldots, g_{2n}$ are analytic functions.

For fixed $\lambda \in \Lambda$ we will denote by $g_\lambda : \mathbb{R}^{n+1} \longrightarrow \mathbb{R}^{2n}$ the mapping defined by $g_\lambda(x) = g(x, \lambda)$.

Let $g_{2n+1}(x) = \omega(x) := x_1^2 + x_2^2 + \ldots + x_{n+1}^2$, and let

$$G_i(x, y, \lambda) := g_i(x, \lambda) - g_i(y, \lambda), \qquad (1 \leqslant i \leqslant 2n)$$
$$G_{2n+1}(x, y) := g_{2n+1}(x) - g_{2n+1}(y) = \|x\|^2 - \|y\|^2,$$

$$G = (G_1, \ldots, G_{2n}) : \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R}^{2n}.$$

Then $G_1, \ldots, G_{2n+1}$ are analytic, and there exist analytic functions h_{ij} such that

$$G_i(x, y, \lambda) = h_{i1}(x, y, \lambda)(x_1 - y_1) + \ldots + h_{i(n+1)}(x, y, \lambda)(x_{n+1} - y_{n+1}).$$

The functions h_{ij} are not uniquely determined.

We fix such functions h_{ij}, and for $1 \leqslant i_1 < i_2 < \ldots < i_{n+1} \leqslant 2n+1$ we define

$$W_{i_1 \ldots i_{n+1}} = \begin{vmatrix} h_{i_1 1} & h_{i_1 2} & \ldots & h_{i_1 (n+1)} \\ h_{i_2 1} & h_{i_2 2} & \ldots & h_{i_2 (n+1)} \\ \vdots & \vdots & \vdots & \vdots \\ h_{i_{n+1} 1} & h_{i_{n+1} 2} & \ldots & h_{i_{n+1} (n+1)} \end{vmatrix}$$

By Cramer's rule

$$(x_1 - y_1) W_{i_1 \ldots i_{n+1}} = \begin{vmatrix} G_{i_1} & h_{i_1 2} & \ldots & h_{i_1 (n+1)} \\ G_{i_2} & h_{i_2 2} & \ldots & h_{i_2 (n+1)} \\ \vdots & \vdots & \vdots & \vdots \\ G_{i_{n+1}} & h_{i_{n+1} 2} & \ldots & h_{i_{n+1} (n+1)} \end{vmatrix}$$

$$\vdots$$

$$(x_{n+1}-y_{n+1})W_{i_1\dots i_{n+1}} = \begin{vmatrix} h_{i_1 1} & h_{i_1 2} & \dots & G_{i_1} \\ h_{i_2 1} & h_{i_2 2} & \dots & G_{i_2} \\ \vdots & \vdots & \vdots & \vdots \\ h_{i_{n+1} 1} & h_{i_{n+1} 2} & \dots & G_{i_{n+1}} \end{vmatrix}$$

Points x and y lie on the same sphere in $\mathbb{R}^{n+1}$ centered at the origin if and only if $\|x\|^2 = \|y\|^2$, i.e. if $G_{2n+1}(x,y) = 0$. If that is the case, then $g(x,\lambda) = g(y,\lambda)$ if and only if $G_1(x,y,\lambda) = \ldots = G_{2n}(x,y,\lambda) = 0$.

Let us define

$$\begin{aligned} A &:= \{(x,y,\lambda) \mid G_1(x,y,\lambda) = \ldots = G_{2n}(x,y,\lambda) = G_{2n+1}(x,y) = 0\} \\ &= \{(x,y,\lambda) \mid \exists_{r>0}\ x,y \in S^n(r),\ g(x,\lambda) = g(y,\lambda)\} \cup \{0\} \times \{0\} \times \Lambda, \\ A_\lambda &:= \{(x,y) \mid (x,y,\lambda) \in A\}. \end{aligned}$$

Then A is closed in $\mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \times \Lambda$.

Let $\Delta := \{(x,x) \mid x \in \mathbb{R}^{n+1}\}$. Then $\Delta \subset A_\lambda$ and $\Delta \times \Lambda$ is closed in $\mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \times \Lambda$. Moreover $(x,y,\lambda) \in \Delta \times \Lambda$ if and only if $x_1 - y_1 = x_2 - y_2 = \ldots = x_{n+1} - y_{n+1} = 0$.

If $(x,y) \in A_\lambda \setminus \Delta$ then $W_{i_1\dots i_{n+1}}(x,y,\lambda) = 0$, for every $1 \leqslant i_1 < \ldots < i_{n+1} \leqslant 2n+1$.

Let us define

$$\begin{aligned} B &:= A \cap \bigcap \{(x,y,\lambda) \mid W_{i_1\dots i_{n+1}}(x,y,\lambda) = 0\}, \\ B_\lambda &:= A_\lambda \cap \bigcap \{(x,y) \mid W_{i_1\dots i_{n+1}}(x,y,\lambda) = 0\} \\ &= \{(x,y) \mid (x,y,\lambda) \in B\}. \end{aligned}$$

Then B is closed in $\mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \times \Lambda$. If $(x,y) \in A_\lambda$ and $x \neq y$, then $(x,y) \in \bigcap \{(x,y) \mid W_{i_1\dots i_{n+1}}(x,y,\lambda) = 0\}$, so

$$A_\lambda \setminus \Delta = B_\lambda \setminus \Delta. \tag{5.1}$$

For $1 \leqslant i, r \leqslant n+1$, if

$$\frac{\partial}{\partial z_i} := \frac{1}{2}\left(\frac{\partial}{\partial x_i} - \frac{\partial}{\partial y_i}\right),$$

then

$$\frac{\partial}{\partial z_i}(x_r - y_r) = \begin{cases} 0, & i \neq r \\ 1, & i = r \end{cases}.$$

For $1 \leqslant j \leqslant 2n+1$,

$$\frac{\partial G_j}{\partial z_i} = \left(\sum_{r=1}^{n+1} \frac{\partial h_{jr}}{\partial z_i}(x_r - y_r) \right) + h_{ji};$$

$$\frac{\partial G_j}{\partial z_i}(x, x, \lambda) = h_{ji}(x, x, \lambda).$$

On the other hand we have

$$\frac{\partial G_j}{\partial z_i}(x, y, \lambda) = \frac{1}{2} \left(\frac{\partial g_j}{\partial x_i}(x, \lambda) + \frac{\partial g_j}{\partial x_i}(y, \lambda) \right);$$
$$\frac{\partial G_j}{\partial z_i}(x, x, \lambda) = \frac{\partial g_j}{\partial x_i}(x, \lambda).$$

Thus

$$h_{ji}(x, x, \lambda) = \frac{\partial g_j}{\partial x_i}(x, \lambda).$$

Lemma 5.1 *Let $\lambda \in \Lambda$. Then $B_\lambda \cap \Delta = \{0\}$ in some neighbourhood of the origin if and only if there exists $r(\lambda) > 0$ such that for all $0 < r < r(\lambda)$ the mapping $g_\lambda = (g_1(\cdot, \lambda), \ldots, g_{2n}(\cdot, \lambda))$ restricted to a sphere of the radius r is an immersion.*

Proof. We have $h_{ji}(x, x, \lambda) = \frac{\partial g_j}{\partial x_i}(x, \lambda)$, so for $(x, x) \in \Delta$ the determinants $W_{i_1 \ldots i_{n+1}}(x, x)$ are the $(n+1) \times (n+1)$–minors of the matrix

$$\begin{bmatrix} \frac{\partial g_1}{\partial x_1} & \frac{\partial g_1}{\partial x_2} & \cdots & \frac{\partial g_1}{\partial x_{n+1}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g_{2n}}{\partial x_1} & \frac{\partial g_{2n}}{\partial x_2} & \cdots & \frac{\partial g_{2n}}{\partial x_{n+1}} \\ \frac{\partial \omega}{\partial x_1} & \frac{\partial \omega}{\partial x_2} & \cdots & \frac{\partial \omega}{\partial x_{n+1}} \end{bmatrix}.$$

The function g_λ satisfies the condition *(b)* of Corollary 4.3 if and only if there exists $r(\lambda) > 0$ such that for $0 < \|x\| < r(\lambda)$ we have $(x, x) \notin B_\lambda$, i.e. $B_\lambda \cap \Delta = \{0\}$ in some neighbourhood of the origin.

□

Let us define

$$\delta : \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \longrightarrow \mathbb{R},$$

$$\delta(x, y) = \|x - y\|^2.$$

Then $\delta \geqslant 0$ and $\delta^{-1}(0) = \Delta$.

Proposition 5.2 *If $\lambda \in \Lambda$ then the following conditions are equivalent:*

(a) there exists $r(\lambda) > 0$ such that for all $0 < r < r(\lambda)$ the mapping g_λ restricted to a sphere of the radius r is an immersion;

(b) $\{0\}$ *is isolated in the set*

$$\Delta \cap \bigcap W_{i_1\ldots i_{n+1}}(\cdot,\lambda)^{-1}(0) = \Delta \cap \bigcap W_{i_1\ldots i_{n+1}}(\cdot,\lambda)^{-1}(0) \cap A_\lambda,$$

where $1 \leqslant i_1 < \ldots < i_{n+1} \leqslant 2n+1$*;*

(c) $B_\lambda \cap \Delta = \{0\}$ *in a neighbourhood of the origin.*

If that is the case, then

(d) in a neighbourhood of the origin $\overline{A_\lambda \setminus \Delta} = B_\lambda$*, and* $(A_\lambda \setminus \Delta) \cup \{0\}$ *is closed;*

(e) if $r > 0$ *is small enough then* $S^n(r/\sqrt{2}) \times S^n(r/\sqrt{2}) \cap (G(\cdot,\cdot,\lambda))^{-1}(0) \setminus \Delta$ *is a compact subset of* $S^n(r/\sqrt{2}) \times S^n(r/\sqrt{2}) \setminus \Delta$.

Proof. Lemma 5.1 implies that $(a) \Leftrightarrow (c)$. Since $\Delta \subset A_\lambda$,

$$\Delta \cap \bigcap_{1\leqslant i_1<\ldots<i_{n+1}\leqslant 2n+1} W_{i_1\ldots i_{n+1}}(\cdot,\lambda)^{-1}(0) \cap A_\lambda = \Delta \cap B_\lambda,$$

so $(b) \Leftrightarrow (c)$. By (5.1) $A_\lambda \setminus \Delta = B_\lambda \setminus \Delta$, so $\overline{A_\lambda \setminus \Delta} = \overline{B_\lambda \setminus \Delta}$. If $B_\lambda \cap \Delta = \{0\}$ in some neighbourhood of the origin, then $\overline{B_\lambda \setminus \Delta} = B_\lambda$ in some neighbourhood of the origin, hence $(c) \Rightarrow (d)$.

By (5.1), $A_\lambda \setminus \Delta = B_\lambda \setminus \Delta$ and by (c) (maybe after making $r(\lambda)$ smaller) for $0 < r < r(\lambda)$ we have $B_\lambda \cap S^{2n+1}(r) \cap \Delta = \emptyset$, so

$$\begin{aligned} B_\lambda \cap S^{2n+1}(r) &= (B_\lambda \setminus \Delta) \cap S^{2n+1}(r) = (A_\lambda \setminus \Delta) \cap S^{2n+1}(r) \\ &= \{(x,y) \in S^{2n+1}(r) \mid \|x\|^2 = \|y\|^2,\ g_\lambda(x) = g_\lambda(y),\ x \neq y\} \\ &= (S^{2n+1}(r) \cap G_{2n+1}^{-1}(0) \cap (G(\cdot,\cdot,\lambda))^{-1}(0)) \setminus \Delta \\ &= S^n(r/\sqrt{2}) \times S^n(r/\sqrt{2}) \cap (G(\cdot,\cdot,\lambda))^{-1}(0)) \setminus \Delta. \end{aligned}$$

B_λ is closed, so $B_\lambda \cap S^{2n+1}(r)$ is compact and we have (e).

□

6 Immersions of spheres — the algebraic case

Assume that $\Lambda \subset \mathbb{R}^p$ is an algebraic set and each g_i is a polynomial. Then there exist polynomials h_{ij} such that

$$G_i(x,y,\lambda) = h_{i1}(x,y,\lambda)(x_1 - y_1) + \ldots + h_{i(n+1)}(x,y,\lambda)(x_{n+1} - y_{n+1}).$$

Now $W_{i_1\ldots i_{n+1}}$ are polynomials, and the sets A, B, A_λ and B_λ are algebraic.

Let us define

$$\begin{aligned} Z &= (\delta^{-1}(0) \times \Lambda) \cap B = (\Delta \times \Lambda) \cap B, \\ Z_\lambda &= \delta^{-1}(0) \cap B_\lambda = \Delta \cap B_\lambda. \end{aligned}$$

So $\delta(x, y, \lambda) := \delta(x, y)$ is continuous on B and the assumptions of [4, Corollary 3.1] hold. Then there exist a finite division $\Lambda = \bigcup S_i$ (where S_i are semialgebraic), continuous, semialgebraic functions $h_i : B \cap (\mathbb{R}^{2n+2} \times S_i) \longrightarrow \mathbb{R}$, and constants $q_i \in \mathbb{Q}^+$ such that for $\lambda \in S_i$, $(x, y) \in B_\lambda$

$$\operatorname{dist}((x, y), Z_\lambda)^{q_i} \leqslant h_i(x, y, \lambda)\delta(x, y).$$

If $\lambda \in S_i$, then there exists such a constant $c_\lambda > 0$ that

$$|h_i(x, y, \lambda)| < c_\lambda$$

for each $(x, y) \in B_\lambda$ sufficiently close to the origin.

Set $d(x, y) = x_1^2 + \ldots + x_{n+1}^2 + y_1^2 + \ldots + y_{n+1}^2 = \|(x, y)\|^2$. From now on we assume that for each $\lambda \in \Lambda$ there exists $r(\lambda) > 0$ such that for each $0 < r < r(\lambda)$ the mapping $g_\lambda | S^n(r)$ is an immersion.

By Proposition 5.2, locally $Z_\lambda = \Delta \cap B_\lambda = \{0\}$. Then for $(x, y) \in B_\lambda$ close to the origin

$$d(x, y)^{\frac{q_i}{2}} = \operatorname{dist}((x, y), \{0\})^{q_i} = \operatorname{dist}((x, y), Z_\lambda)^{q_i} \leqslant c_\lambda \delta(x, y).$$

If $\alpha > \max_i\{\frac{q_i}{2}\}$ is an integer, then for each $\lambda \in \Lambda$ and $t > 0$

$$\delta(x, y) \geqslant t d(x, y)^\alpha \tag{6.1}$$

for $(x, y) \in B_\lambda$ lying sufficiently close to the origin. By (5.1) $A_\lambda \setminus \Delta = B_\lambda \setminus \Delta$, so the inequality (6.1) holds on $A_\lambda \setminus \Delta$, i.e. on

$$\{(x, y) \mid G_1(x, y, \lambda) = \ldots = G_{2n}(x, y, \lambda) = \|x\|^2 - \|y\|^2 = 0\} \setminus \delta^{-1}(0).$$

By Proposition 5.2 (d), $(A_\lambda \setminus \Delta) \cup \{0\}$ is closed.

A polynomial $F_1(x, y) = \|x\|^2 - \|y\|^2$ has an isolated critical point at the origin. Then

$$L(r) = S^{2n-1}(r) \cap F_1^{-1}(0) = S^n(r/\sqrt{2}) \times S^n(r/\sqrt{2})$$

is a compact oriented $2n$–dimensional manifold. Since $\delta(x, y) = \|x - y\|^2 \geqslant 0$,

$$U(r) = L(r) \setminus \delta^{-1}(0) = S^n(r/\sqrt{2}) \times S^n(r/\sqrt{2}) \setminus \Delta.$$

By Proposition 2.2 there exists $\alpha_0 > 0$ such that for any integer $\alpha > \alpha_0$ and $t \neq 0$, $(F_1, \delta - t d^\alpha)^{-1}(0)$ has an isolated singularity at the origin. (As F_1, δ and d are homogeneous of degree 2, it is easy to verify that $\alpha_0 = 1$ would be enough.)

Let us assume that n is even. Take an integer $\alpha > \max(\max_i\{\frac{q_i}{2}\}, \alpha_0)$. Put $F_2 = \delta - t d^\alpha$ for arbitrary $t > 0$, and

$$F = (F_1, F_2),$$
$$G_\lambda = G(\cdot, \cdot, \lambda) : \mathbb{R}^{n+1} \times \mathbb{R}^{n+1}, 0 \longrightarrow \mathbb{R}^{2n}, 0.$$

By Proposition 5.2 (e), $L(r) \cap G_\lambda^{-1}(0) \setminus \delta^{-1}(0)$ is a compact subset of $U(r)$. By Proposition 2.3, for $r > 0$ small enough, the topological degree of

$$(S^n(\tfrac{r}{\sqrt{2}}) \times S^n(\tfrac{r}{\sqrt{2}}) \setminus \Delta, S^n(\tfrac{r}{\sqrt{2}}) \times S^n(\tfrac{r}{\sqrt{2}}) \setminus (G_\lambda^{-1}(0) \cup \Delta)) \longrightarrow (\mathbb{R}^{2n}, \mathbb{R}^{2n} \setminus \{0\})$$

$$(x, y) \mapsto G_\lambda(x, y) = g_\lambda(x) - g_\lambda(y),$$

i.e. $2\,\mathrm{I}(g_\lambda | S^n(r/\sqrt{2}))$, equals the local topological degree $\deg_0(F, G_\lambda)$.

Let us define a polynomial mapping $H : \mathbb{R} \times \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}^{2n}$:

$$H(t, x, y, \lambda) = (F_1(x, y), \delta(x, y) - td(x, y)^\alpha, g(x, \lambda) - g(y, \lambda)).$$

Denote $H_\lambda^t(x, y) = H(t, x, y, \lambda)$. For any $\lambda \in \Lambda$ and $t > 0$, the mapping H_λ^t has an isolated zero at the origin and the local topological degree $\deg_0(H_\lambda^t) = 2\,\mathrm{I}(g_\lambda | S^n(r))$ for $0 < r < r(\lambda)$. If $t < 0$ then $\delta - td^\alpha > 0$, except of the origin, so that $\deg_0(H_\lambda^t) = 0$.

Theorem 6.1 *Let $\Lambda \subset \mathbb{R}^p$ be an algebraic set and let n be an even integer. Let*

$$g_1(x, \lambda), \ldots, g_{2n}(x, \lambda) : \mathbb{R}^{n+1} \times \Lambda \longrightarrow \mathbb{R}^{2n}$$

be polynomials. Assume that for each $\lambda \in \Lambda$ there exists $r(\lambda) > 0$ such that for each $0 < r < r(\lambda)$ the mapping g_λ restricted to a sphere $S^n(r)$ of the radius r centered at the origin is an immersion. Then the function $\Lambda \ni \lambda \mapsto \mathrm{I}(g_\lambda | S^n(r)) \in \mathbb{Z}$ is algebraically constructible.

Proof. Using the same arguments as in [9, Theorem 3.4] one may prove that there exist polynomials $h_1, \ldots, h_s$ on $\mathbb{R} \times \Lambda$, such that if 0 is isolated in $(H_\lambda^t)^{-1}(0)$ then

$$\deg_0(H_\lambda^t) = \sum_{i=1}^{s} \operatorname{sgn} h_i(t, \lambda),$$

i.e. this function is algebraically constructible.

For $r > 0$ small enough

$$\begin{aligned} \mathrm{I}(g_\lambda | S^n(r)) &= \lim_{t \to 0} \frac{1}{2} (\deg_0(H_\lambda^t) + \deg_0(H_\lambda^{-t})) \\ &= \lim_{t \to 0} \frac{1}{2} \sum_{i=1}^{s} (\operatorname{sgn} h_i(\lambda, t) + \operatorname{sgn} h_i(\lambda, -t)) =: \psi(\lambda). \end{aligned}$$

According to [9, Lemma 6.5] the function $\Lambda \ni \lambda \mapsto \psi(\lambda) \in \mathbb{Z}$ is algebraically constructible, and so is $\Lambda \ni \lambda \mapsto \mathrm{I}(g_\lambda | S^n(r))$.

□

Remark. It is easy to see that if mappings g_λ are immersions on small spheres only for $\lambda \in \Lambda \setminus \Sigma$, where Σ is a proper algebraic subset of Λ, then the function $\Lambda \ni \lambda \mapsto \mathrm{I}(g_\lambda | S^n(r)) \in \mathbb{Z}$ is generically algebraically constructible, i.e. it coincides with an algebraically constructible function in $\Lambda \setminus \Sigma$.

References

[1] Akbulut, S., King, H.: Topology of real algebraic sets. MSRI Publ. **25**, New York: Springer – Verlag, 1992

[2] Coste, M.: Real Algebraic Sets, Summer School and Conference on Real Algebraic Geometry and its Applications (2003)

[3] Coste, M., Kurdyka, K.: Le discriminant d'un morphisme de variétés algébriques réelles. *Topology* **37** no. 2, 393–399 (1998)

[4] Fekak, A.: Exposants de Łojasiewicz pour les fonctions semi–algébriques. *Annales Polonici Mathematici* **56** 123–131 (1992)

[5] Lashof, R., Smale, S.: On the immersion of manifolds in euclidean space. *Ann. of Math.* (2) **68**, 562–583 (1958)

[6] McCrory, C., Parusiński, A.: Algebraically constructible functions. *Ann. Sci. École Norm. Sup.* (4) **30**, no.4, 527–552 (1997)

[7] McCrory, C., Parusiński, A.: Topology of real algebraic sets of dimension 4: necessary conditions. *Topology* **39** no. 3, 495–523 (2000)

[8] McCrory, C., Parusiński, A.: Algebraically Constructible Functions: Real Algebra and Topology, preprint (2002)

[9] Parusiński, A., Szafraniec, Z.: Algebraically constructible functions and signs of polynomials. *Manuscripta Math.* **93**, no. 4, 443–456 (1997)

[10] Parusiński, A., Szafraniec, Z.: On the Euler characteristic of fibres of real polynomial maps. Singularities Symposium—Łojasiewicz 70 (Kraków, 1996; Warsaw, 1996), *Banach Center Publ.* **44** Polish Acad. Sci., Warsaw (1998), 175–182 (1998)

[11] Smale, S.: The classification of immersions of spheres in Euclidean spaces. *Ann. of Math.* (2) **69**, 327–344 (1959)

[12] Szafraniec, Z.: On the Euler characteristic of analytic and algebraic sets. *Topology* **25**, 411–414 (1986)

[13] Whitney, H.: The self–intersections of a smooth n–manifold in $2n$–space. *Ann. of Math.* (2) **45**, 220–246 (1944)

[14] Whitney, H.: Differentiable manifolds. *Ann. of Math.* (2) **37**, no.3, 645–680 (1936)